

Climate change and polar range expansions: Could cuttlefish cross the Arctic?

José C. Xavier^{1,2}, Lloyd S. Peck¹, Peter Fretwell¹ & John Turner¹

1- British Antarctic Survey, Natural Environment Research Council, High Cross, Madingley Road, CB3 0ET Cambridge, UK

2- Marine and Environmental Science Centre (MARE), Faculdade de Ciências e Tecnologia, Universidade de Coimbra, Departamento das Ciências da Vida, Coimbra, PT

Abstract

Climate change can have major effects on the distribution of species. In marine ecosystems, the cold waters of the Arctic have restricted warmer water species from crossing between Eurasia and North America. However, with Arctic waters becoming warmer, various marine species have expanded their distribution. Cuttlefish are fast growing, voracious predators and are absent in American waters. The European cuttlefish *Sepia officinalis* is the most northerly-distributed cuttlefish, with potential to expand its range and cross to the American continent, potentially causing changes in shelf food webs. Climate model predictions suggest that the *S. officinalis* could potentially reach American shores, by 2300 via the north Atlantic with medium mitigation of greenhouse gas concentrations; we predict that adult dispersal of cuttlefish across the Atlantic sector would require a migration distance of over 1400 km at depths below 200 m and temperatures above 7°C (temperature below which cuttlefish can not maintain routine metabolic processes physiologically). For temperatures above 9.5°C (temperature above which cuttlefish can grow), 2500 km would be required, and such conditions will possibly exist by the year 2300. If they reach American shores they could have large impacts on coastal marine ecosystems, due to their wide diet (e.g. diet covers many shallow water crustacean and fish species) and its potential as prey, and due to their short life history strategy of “live fast, die young”.

KEYWORDS: Invasion, Polar, predator, cephalopod, distribution - Climate change - *Sepia officinalis*

34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61

INTRODUCTION

Global warming is producing significant changes in marine ecosystems. Emerging evidence suggests that marine organism distributions may respond more rapidly to climate change than those on land, despite slower oceanic warming (Richardson and Poloczanska 2008; Kortsch et al. 2015). Range shifts of hundreds of kilometres in a few decades have been observed in various organisms, such as phytoplankton, zooplankton and fish (Beaugrand et al. 2002). However, studies have rarely investigated fast growing, carnivorous species with the potential for also marked impacts on new habitats. The short generation times of coastal cuttlefish, squid and octopods might enable them to adjust faster to changing environmental conditions than those with slow growth and late maturity (Boyle and Rodhouse 2005; Xavier et al. 2015). Indeed, the short life span and generation time of shallow water cephalopods has been suggested to enable a proportion of each new generation to actively avoid or exploit localised warming events (Rodhouse 2013). Some studies already investigated range shifting in Octopods, such as *Octopus tetricus* (Ramos et al. 2014) at the southerly extension of the warm East Australian Current, showing that this octopod maintains a fast growth rate (under warming conditions), high rate of population turnover, small body size and a short life span at the leading edge of their range extension. In squid, such as *Dosidicus gigas* (Zeidberg and Robison 2007; Ruiz-Cooley et al. 2013), *Sepietta oweniana*, *Todarodes sagittatus*, *Todaropsis eblanae* and *Teuthowenia megalops* (Golikov et al. 2013; Golikov et al. 2014), range expansion has been attributed to changes in climate-linked oceanographic conditions (e.g. warming waters), a reduction in competing top predators and/or of a decline in abundance of a commercial groundfish species. However, unlike octopods and squid, cuttlefish *Sepia* spp. do not occur in the Americas or Polar regions (Boletzky 1983 ; Xavier et al. 1999; Xavier and Cherel 2009; Rodhouse et al. 2014). Nearly 115 sepiid species are described inhabiting shallow tropical/temperate waters of Afro-Eurasia, along the coasts of the East Atlantic, Indian and West Pacific (Reid et al. 2005). The question arises: as the Arctic warms when will environmental conditions allow cuttlefish to cross the Arctic and reach American coastal waters?

FACTORS AFFECTING CUTTLEFISH DISTRIBUTION

Here, we focus our discussion on the European cuttlefish, *S.officinalis*, and the potential range expansion of the species to America via Europe. The route via the Bering Sea (i.e. Asia - America route)

65 is not discussed further in this study because there are no published studies on the temperature tolerance
66 of *Sepia kabiensis* (the most northern cuttlefish species distributed in the north Pacific) (Reid et al. 2005)
67 nor any other North Pacific, cold-adapted cuttlefish species to support such hypotheses. However, due to
68 the shallow waters between Asia and the Americas, via the Bering Sea, the dominating W-E currents
69 could aid such range expansion of cuttlefish.

70 The most northerly distributed cuttlefish worldwide is *Sepia officinalis* (Family Sepiidae, Order
71 Sepiida, Class Cephalopoda) at around 62°N in European waters, a well known opportunistic and
72 voracious marine predator that can reach large size (up to 4-kg), and feeds on a wide range of prey (Reid
73 et al. 2005). Physiologically, *S. officinalis* cannot maintain routine metabolic processes at temperatures
74 $\leq 7^{\circ}\text{C}$, grows positively from water temperatures above 9.5°C and spawning peaks at $13 - 15^{\circ}\text{C}$ (Boletzky
75 1983 ; Pimentel et al. 2012). Temperature has large effects on development in this species. When eggs
76 are exposed to warming conditions (13 to 19°C), oxygen consumption increases throughout
77 embryogenesis. At 15°C , 41% of the egg yolk is used for growth, and only 10% is used for catabolic
78 processes, whereas at higher temperatures (24°C) yolk utilisation for growth is only 15%, but 52% is for
79 catabolism, resulting in small hatchling size (Bouchaud 1991). *S. officinalis* is thus well adapted to cool
80 temperate conditions and it has been suggested that, with warming, *S. officinalis* may lay their eggs in
81 warmer and deeper environments (Pimentel et al. 2012). *S. officinalis* is, however, limited to a depth of
82 150-200 m as the chambers in the cuttlebone, the calcified internal shell, can implode beyond these
83 depths (Ward and Boletzky 1984). Other cuttlefish species at lower latitude and warmer sites, on the other
84 hand, can be found up to 1000 m deep, and may be better adapted to pressure. *Sepia elegans*, for example
85 is found up to 60°N in depths of 500 m, and *Sepia orbignyana* up to 54°N in depths of 570 m (Reid et al.
86 2005).

87 Cuttlefish today are absent from the Americas, but this is probably not due to lack of habitat (Young
88 et al. 1998), as potential prey are abundant. In fact, *Sepia officinalis* has been cultured during multiple
89 generations in American laboratories where they have been fed with live local crustaceans and fish
90 (Forsythe et al. 2002). One reason for their absence, appears to be low temperature in the Arctic and the
91 depth/distance barriers of the other ocean basins, combined with the time of formation of these barriers
92 (Young et al. 1998). The shallow-water tropical bridge between Africa and South America was severed in
93 the late Cretaceous, leaving the northern rim of the Atlantic Basin as the only possible migration route for
94 shallow demersal organisms between Europe and the Americas after this time. During the early Cenozoic
95 (Palaeocene and Eocene), a series of radiations began in the warm Tethys Sea that resulted in the broad
96 colonization of the Belosaepiidae (a cephalopod family known from the Eocene, closely similar to
97 cuttlefish) (Khromov 1998). Migration across the Atlantic was possible around the top of the basin,

98 where temperatures were tolerable. After the Belosaepiidae became extinct in the Oligocene, cuttlefish
99 species emerged in Europe (North East Atlantic), the only location of fossils currently known (Young et
100 al. 1998). Their distribution was, however, restricted by the colder conditions during this period,
101 preventing colonisation across the North Atlantic as utilised by the Belosaepiidae.

102 Cuttlefish are also absent from the Antarctic, which is likely due to cold conditions and the deep water
103 barriers between Antarctica and the other land masses, that were completed with the formation of the
104 Drake Passage around 30-50 million years (Ma) ago (Livermore et al. 2005). There are cephalopod fossils
105 in the Antarctic, from belemnites and Mesozoic teuthids (Doyle 1991), but none from the family Sepiidae
106 probably due to the absence of marine Tertiary sediments (Dirk Fuchs, Freie Universität Berlin, pers.
107 comm.). The sepiids closest to the Antarctic are in Australian waters, at 42°S, but a lack suitable habitat
108 further south (Reid et al. 2005).

109 The northern expansion of *S. officinalis* appears to be mainly limited by physiology, but life cycle
110 constraints and habitat availability may also play a part. Eggs of this species, as for all sepiids, are large
111 and attached to hard substrata in depths usually less than 50 m, producing benthic hatchlings with limited
112 dispersal capacity. Lifespan is 18-24 months and dispersal is predominantly via sub-adult and adult
113 migration (Reid et al. 2005). As cuttlefish spawning appears restricted to shallow depths, iceberg scouring
114 might also limit their ability to colonise low temperature areas by restricting areas of suitable habitat to
115 attach eggs to algae and hard substrata. If waters warmed, *S. officinalis* could potentially expand its
116 northern range edge via two routes: either along the 200 m shelf towards the Arctic, and/or via the Faroe
117 Islands – Iceland - Greenland to north Canada over deeper waters (Fig. 2), both aided by surface currents
118 (Straneo and Heimbach 2013) that could transport adult cuttlefish (as they expand north) and/or, for
119 example, rafting with marine (floating) debris (e.g. broken kelp, floating old fishing gear or plastic
120 material initially at the bottom) (as cuttlefish eggs) (Boletzky 1983; Arkley et al. 1996; Blanc and
121 Daguzan 1998; Sykes et al. 2014). Although cuttlefish are benthopelagic organisms, they are known to
122 also swim into the water column (Okutani 1990), which could be transport them further north by currents.
123 Furthermore, if these adults can reach Faroe Islands, Iceland, Greenland and North Canada, they would be
124 able to reproduce and attach their eggs to available algae (e.g. sea grass or kelp) (Kjellman 1883; Mann
125 1973; Arkley et al. 1996; Short et al. 2007). This would permit a sequence of generations to allow the
126 establishment of cuttlefish in these regions.

127 Depth may not be a total barrier for cuttlefish expansion, as between South Africa and Mauritius
128 (Mascarene Ridge), the same cuttlefish species (*Sepia vermiculata*) occurs in both places, although
129 separated by deep waters (Reid et al. 2005). If cuttlefish cross the Arctic, their life history characteristics

would potentially allow them to adapt quickly and spread rapidly. A further consideration is that the aquarium industry might provide another colonisation or invasion route, as cuttlefish are readily available in the USA as pets. If released in the sea, under the right conditions, they could flourish, as has occurred for other invaders (Gido and Brown 1999).

PREDICTING THE NORTHERN RANGE EXPANSION OF *S. OFFICINALIS*

How much do Arctic waters need to warm for cuttlefish to cross the Arctic? As *S. officinalis* is the most abundant, largest and the most known northerly distributed cuttlefish, we use this species as the candidate for exploring the likelihood of range expansion. The total reported annual landings of cuttlefish in Europe range from 35 000 to 41 000 tons, showing that a considerable population exists in European waters (Pierce et al. 2010). Recent Arctic Ocean warming is thought to be closely connected with increased heat content of the Atlantic water masses with a 1.3°C increase in annual water temperatures from 1990 to 2005 (Walczowski and Piechura 2006).

An expected migration route for *S. officinalis* across the Arctic would be from northern Europe via Iceland to North America (Fig. 1). Based on the HadGEM2-ES model (Hadley Centre Global Environmental Model version 2 Earth System configuration) with medium mitigation of greenhouse gas concentrations (following the “Representative Concentration Pathway” RCP4.5) (Caesar et al. 2013), we predict that adult dispersal of cuttlefish across the Atlantic sector would require a migration of over 1400 km at depths deeper than 200 m and temperatures above 7°C. For temperatures above 9.5°C, 2500 km would be required, and such conditions will possibly exist by the year 2300 (See supplementary material).

Increasing temperatures may facilitate a range expansion of cuttlefish to the Arctic, but climate change may also bring environmental challenges such as ocean acidification. Polar marine calcifying organisms may be amongst the first affected by ocean acidification-driven changes in marine carbonate chemistry (Orr et al. 2005). In cuttlefish laboratory experiments with *S. officinalis* reported higher cuttlebone calcification associated with decreasing pH (Dorey et al. 2013). This hyper-calcification would likely change the cuttlebone buoyancy, and possibly change the implosion resistance of this structure. However, the behavioural or ecological significance of these changes are still unknown. When compared with other organisms, Perry et al. (2005) showed that the distributions of both exploited and non-exploited fish species have responded markedly to recent increases in sea temperature, with nearly two-thirds of species shifting in mean latitude or depth over 25 years. Furthermore, most of the fish species that shifted their distributions had faster life cycles (but still slower than cuttlefish) and smaller body

sizes. Temperature rises are likely to have profound impacts on community interactions through continued shifts in distribution of marine organisms (Perry et al. 2005), particularly those that are fast growing, with short life cycles and a broad range of prey.

In summary, the cuttlefish *Sepia officinalis* is one of the marine organisms that, under a favourable climatic scenario and conditions (e.g. by currents, cross over deep regions), may be able to cross the Arctic region and occupy a niche in the marine ecosystems of the American continent. Due to its fast growth and metabolism, typical of cephalopods, cuttlefish may be able to occupy an ecological niche rapidly.

ACKNOWLEDGEMENTS

We thank Roger Villanueva for the numerous discussions and contributions to previous drafts and Tom Bracegirdle, Dan Jones, Alexey Golikov and Emma Boland for advice on polar oceanography. This research is part of the SCAR AnT-ERA and ICED programs. JX is supported by the Investigator FCT program (IF/00616/2013).

References

- Arkley K, Jacklin MS, Boulter M, Tower J (1996) The cuttlefish (*Sepia officinalis*): A guide to its exploitation in UK waters, Seafish Report N. SR467. Sea-fish Industry Authority, Hull, UK
- Beaugrand G, Reid PC, Ibañez F, Lindley JA, Edwards M (2002) Reorganization of North Atlantic marine copepod biodiversity and climate. *Science* 296: 1692
- Blanc A, Daguzan J (1998) Artificial surfaces for cuttlefish eggs (*Sepia officinalis* L.) in Morbihan Bay, France. *Fish Res* 38: 225-231
- Boletzky SV (1983) *Sepia officinalis*. In: Boyle PR (ed) Cephalopod life Cycles. Academic Press, London, pp 31-52
- Bouchaud O (1991) Energy consumption of the cuttlefish *Sepia officinalis* (Mollusca: Cephalopoda) during embryonic development, preliminary results. *Bull of Mar Sci* 49: 333-340
- Boyle P, Rodhouse PG (2005) Cephalopods ecology and fisheries. Blackell Science, Oxford
- Caesar J, Palin E, Liddicoat S, Lowe J, Burke E, Pardaens A, Sanderson M, Kahana R (2013) Response of the HadGEM2 Earth System Model to Future Greenhouse Gas Emissions Pathways to the Year 2300. *J Climate* 26: 3275–3284
- Dorey N, Melzner F, Martin S, Oberhänsli F, Teyssié JL, Bustamante P, Gattuso JP, Lacoue-Labarthe T (2013) Ocean acidification and temperature rise: effects on calcification during early development of the cuttlefish *Sepia officinalis*. *Mar Biol* 160: 2007-2022
- Doyle P (1991) Teuthid cephalopods from the upper Jurassic of Antarctica. *Palaentology* 34: 169-178
- Forsythe J, Lee P, Walsh L, Clark T (2002) The effects of crowding on growth of the European cuttlefish, *Sepia officinalis* Linnaeus, 1758 reared at two temperatures. *J Exp Mar Biol and Ecol* 269: 173-185
- Gido KB, Brown JH (1999) Invasion of North American drainages by alien fish species. *Freshw Biol* 42: 387-399
- Golikov AV, Sabirov RM, Lubin PA, Jørgensen LL (2013) Changes in distribution and range structure of Arctic cephalopods due to climatic changes of the last decades. *Biodiversity* 14: 24-35
- Golikov AV, Sabirov RM, Lubin PA, Jørgensen LL, Beck I-M (2014) The northernmost record of *Sepietta oweniana* (Cephalopoda: Sepiolidae) and comments on boreo-subtropical cephalopod species occurrence in the Arctic. *Mar Biodiversity Records* 7: e58

202 Khromov DN (1998) Distribution patterns of Sepiidae. In: Voss NA, Vecchione M, Toll RB, Sweeney MJ (eds)
 203 Systematics and biogeography of cephalopods. Smithsonian Contr Zool, pp 191-206
 204 Kjellman FR (1883) The algae of the Arctic Sea: A survey of the species, together with an exposition of the general
 205 characters and the development of flora. P. A. Norstedt and Soner, Stockholm
 206 Kortsch S, Primicerio R, Fossheim M, Dolgov AV, Aschan M (2015) Climate change alters the structure of arctic
 207 marine food webs due to poleward shifts of boreal generalists. Proc R Soc B 282: 20151546
 208 Livermore R, Nankivell A, Eagles G, Morris P (2005) Paleogene opening of Drake Passage. Earth and Planetary
 209 Letters 236: 459-470
 210 Mann KH (1973) Seaweeds: their productivity and strategy for growth. Science 182: 975-981
 211 Okutani T (1990) Squids, cuttlefish and octopuses. Mar Behav and Physiol 18: 1-17 doi
 212 10.1080/10236249009378778
 213 Orr JC, Fabry VJ, Aumont O, Bopp L, Doney SC, Feely RA, Gnanadesikan A, Gruber N, Ishida A, Joos F, Key RM,
 214 Lindsay K, Maier-Reimer E, Matear R, Monfray P, Mouchet A, Najjar RG, Plattner G-K, Rodgers KB, Sabine
 215 CL, Sarmiento JL, Schlitzer R, Slater RD, Totterdell IJ, Weirig M-F, Yamanaka Y, Yool A (2005)
 216 Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms.
 217 Nature 437: 681-686
 218 Perry AL, Low PJ, Ellis JR, Reynolds JD (2005) Climate Change and Distribution Shifts in Marine Fishes. Science 308:
 219 1912-1915
 220 Pierce GJ, Allcock AL, Bruno I, Bustamante P, González A. F, Guerra A, Jereb P, Lefkaditou E, Malham SK, Moreno
 221 A, Pereira J, Piatkowski U, Rasero M, Sánchez P, Santos MB, Santurtun M, Seixas S, Sobrino I, Villanueva R
 222 (2010) Cephalopod biology and fisheries in Europe. International Council for the Exploration of the Sea,
 223 Copenhagen
 224 Pimentel MS, Trubenbach K, Faleiro F, Boavida-Portugal J, Repolho T, Rosa R (2012) Impact of ocean warming on
 225 the early ontogeny of cephalopods: a metabolic approach. Mar Biol 159: 2051-2059
 226 Ramos JE, Pecl GT, Moltschaniwskyj NA, Strugnell JM, León RI, Semmens JM (2014) Body size, growth and life
 227 span: implications for the polewards range shift of *Octopus tetricus* in south-eastern Australia. PLoS ONE
 228 9: e103480
 229 Reid A, Jereb P, Roper CFE (2005) Family Sepiidae. In: Jereb P, Roper CFE (eds) Cephalopods of the world An
 230 annotated and illustrated catalogue of species known to date. FAO Species Catalogue for Fishery
 231 Purposes. Rome, FAO, pp 57-152
 232 Richardson AJ, Poloczanska ES (2008) Ocean Science: under-resourced, under threat. Science 320: 1294-1295
 233 Rodhouse PG (2013) Role of squid in the Southern Ocean pelagic ecosystem and the possible consequences of
 234 climate change. Deep Sea Res II 95: 129-138
 235 Rodhouse PG, Xavier JC, Griffiths H (2014) Southern Ocean squid. In: De Broyer C, Koubbi P, Griffiths H, Danis B,
 236 David B, Grant S, Gutt J, Held C, Hosie G, Huettmann F, Post A, Raymond B, Robert-Coudert Y, van de
 237 Putte A (eds) The CAML / SCAR-MarBIN Biogeographic Atlas of the Southern Ocean. Scientific Committee
 238 on Antarctic Research, Cambridge, pp 284-289
 239 Ruiz-Cooley RI, Ballance LT, McCarthy MD (2013) Range expansion of the jumbo squid in the NE Pacific: $\delta^{15}\text{N}$
 240 decrypts multiple origins, migration and habitat use. PloS one 8: e59651
 241 Short F, Carruthers T, Dennison W, Waycott M (2007) Global seagrass distribution and diversity: A bioregional
 242 model. J Exp Mar Biol and Ecol 350: 3-20 doi <http://dx.doi.org/10.1016/j.jembe.2007.06.012>
 243 Straneo F, Heimbach P (2013) North Atlantic warming and the retreat of Greenland's outlet glaciers. Nature 504:
 244 36-43
 245 Sykes AV, Domingues P, Andrade JP (2014) *Sepia officinalis*. In: Iglesias J, Fuentes L, Villanueva R (eds) Cephalopod
 246 Culture. Springer, pp 175-204
 247 Walczowski W, Piechura J (2006) New evidence of warming propagating toward the Arctic Ocean. Geophys Res
 248 Lett 33: L12601
 249 Ward P, Boletzky SV (1984) Shell implosion depth and implosion morphologies in three species of *Sepia*
 250 (Cephalopoda) from the Mediterranean Sea. J Mar Biol Ass UK 64: 955-966

251 Xavier JC, Allcock L, Cherel Y, Lipinski MR, Gomes-Pereira JN, Pierce G, Rodhouse PGK, Rosa R, Shea L, Strugnell J,
 252 Vidal E, Villanueva R, Ziegler A (2015) Future challenges in cephalopod research. J Mar Biol Ass UK 95:
 253 999-1015
 254 Xavier JC, Cherel Y (2009) Cephalopod beak guide for the Southern Ocean. British Antarctic Survey
 255 Xavier JC, Rodhouse PG, Trathan PN, Wood AG (1999) A Geographical Information System (GIS) atlas of
 256 cephalopod distribution in the Southern Ocean. Antarct Sci 11: 61-62
 257 Young RE, Vecchione M, Donovan D (1998) The evolution of coleoid cephalopods and their present biodiversity
 258 and ecology. S Afr J Mar Sci 20 393-420
 259 Zeidberg LD, Robison BH (2007) Invasive range expansion by the Humboldt squid, *Dosidicus gigas*, in the eastern
 260 North Pacific. Proc Nat Acad Sci 104: 12948-12950

261
 262


1 SUPPLEMENTARY MATERIAL. Distances across the Arctic from Europe to North America (upper)
2 following the mean summer sea surface temperature contours of 7 °C (limit for *S. officinalis* adult
3 survival) and 9.5°C (limit for *S. officinalis* reproduction) as predicted by runs of the HadGEM2-
4 ES model using a medium mitigation run RCP4.5 (See methods in Caesar *et al.* 2013).

Europe to North America
(via Iceland)


	7 °C	7 °C	9.5 °C	9.5 °C
Year	Distance (km) coast to coast	Distance (km) to reach the 200 metre contour depth in North America East coast	Distance (km) coast to coast	Distance (km) to reach the 200 metre contour depth in North America East coast
2012	2960	2530	3042	2530
2020	2960	2530	3023	2530
2030	2953	2530	2953	2530
2040	2953	2530	2953	2530
2050	2953	2530	2953	2530
2060	2953	2530	2953	2530
2070	2953	2439	2953	2530
2080	2953	2439	2953	2530
2090	2930	2388	2953	2530
2100	2930	2064	2953	2530
2110	2930	2062	2953	2530
2120	2930	2008	2953	2530
2130	2930	2008	2953	2530
2140	2930	2008	2953	2530
2150	2930	2008	2953	2530
2160	2930	2008	2953	2530
2170	2930	2008	2953	2530
2180	2930	2008	2953	2530
2190	2930	2008	2953	2530
2200	2930	2008	2953	2530
2210	2930	1932	2953	2530
2220	2930	1781	2953	2530
2230	2930	1718	2953	2530
2240	2930	1487	2953	2530
2250	2930	1487	2953	2530
2260	2930	1469	2953	2530
2270	2930	1430	2953	2530
2280	2930	1430	2953	2530
2290	2930	1430	2953	2530
2300	2930	1430	2953	2530

5
6
7

Present day SST



Projected SST for 2300



SST

C



Figure 1. Mean summer sea surface temperatures (SST in °C) for the present day and projected for 2300, for the North Atlantic (The “present SST” also includes the present distribution of *Sepia officinalis*; Eastern Atlantic, from the Shetland Islands and southern Norway south through the Mediterranean Sea to northwestern Africa up to Senegal (Reid et al. 2005)). Projections are based from runs of the HadGEM2-ES medium mitigation run (see text) and are for mean summer temperature covering a 3 month period. The 200 m depth contours is also shown in all panels.




Figure 2. Predicted best migration routes of *Sepia officinalis* for the North Atlantic projected for 2300, under a mean summer sea surface temperatures (SST in °C) obtained from HadGEM2-ES medium mitigation run: solid arrows for optional route with water temperatures above 7°C and dash lines for the route above 9°C. The 200 m contour is also shown. See figure 1 for water temperatures for the corresponding colours.